

UNIVERSITÄTSKLINIKUM HAMBURG-EPPENDORF

Institut und Poliklinik für Medizinische Psychologie

Direktor: Prof. Dr. med. Dr. phil. Martin Härter

Incomplete Medication Adherence of Chronically Ill Patients in German Primary Care

Dissertation

zur Erlangung des Grades eines Doktors der Medizin
an der Medizinischen Fakultät der Universität Hamburg.

vorgelegt von:

Jakob Hüther
aus Freiburg im Breisgau

Hamburg 2013

**Angenommen von der
Medizinischen Fakultät der Universität Hamburg am: 27.05.2014**

**Veröffentlicht mit Genehmigung der
Medizinischen Fakultät der Universität Hamburg.**

Prüfungsausschuss, der/die Vorsitzende: Prof. Dr. M. Härter

Prüfungsausschuss, zweite/r Gutachter/in: Prof. Dr. H. v. d. Bussche

CONTENT

ARTICLE.....	4
SUMMARY.....	11
AUTHOR CONTRIBUTION	24
ACKNOWLEDGEMENTS.....	25
CURRICULUM VITAE	26
AFFIDAVIT (EIDESSTATTLICHE VERSICHERUNG)	27

Incomplete medication adherence of chronically ill patients in German primary care

Jakob Hüther¹
 Alessa von Wolff¹
 Dorit Stange²
 Martin Härter¹
 Michael Baehr²
 Dorothee C Dartsch³
 Levente Kriston¹

¹Department of Medical Psychology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany;

²Hospital Pharmacy, University Medical Center Hamburg-Eppendorf, Hamburg, Germany; ³Institute of Pharmacy, University of Hamburg, Hamburg, Germany

This article was published in the following Dove Press journal:

Patient Preference and Adherence

21 March 2013

Number of times this article has been viewed

Background: Incomplete medication adherence is a major problem in health care worldwide. Patients who adhere to medical treatment have a better prognosis and create fewer costs.

Objective: To assess the degree of incomplete adherence of chronically ill routine primary care patients in a German setting and analyze the association between incomplete medication adherence, as well as clinical and sociodemographic patient characteristics.

Methods: In a cross-sectional survey, chronically ill patients were asked to assess their adherence in primary care retrospectively using the Medication Adherence Report Scale (MARS-D) questionnaire. To investigate the association of incomplete adherence with sociodemographic and clinical data, univariate and multivariate analyses were conducted.

Results: In total, 62.1% of 190 patients were categorized as incompletely adherent. The mean MARS-D score was 23.5 (standard deviation = 2.7). Analyses revealed no statistically significant associations at $P < 0.05$ between degree of adherence and patient characteristics. The total explained variance amounted to 11.8% (Nagelkerke's $R^2 = 0.118$) in the multivariate analysis.

Conclusion: Previously reported results regarding associations of sociodemographic and clinical data with incomplete medication adherence could not be confirmed for this sample of chronically ill patients. In order to be able to provide guidelines for the reduction of incomplete medication adherence in German primary care, further research is needed.

Keywords: medication, adherence, chronic illness, primary care, Germany, MARS

Introduction

In all health care systems, prescribing medication represents a major part of medical treatment. Medications have been shown to improve health outcomes and reduce the utilization of health care resources.¹⁻⁵ Moreover, it has been shown that patients who are less likely to take their prescribed medication are more likely to suffer from secondary diseases,⁶ and thus to be hospitalized or die.⁷⁻¹⁰ The World Health Organization defines adherence as the extent to which a person's behavior (eg, taking medication, following a diet, and/or executing lifestyle changes) corresponds with agreed recommendations from a health care provider.¹¹ Therefore, incomplete adherence is the occurrence of a patient not completely following these recommendations.

Beyond the negative effects of incomplete adherence on a patient's health, it also represents a financial burden on health care systems. The estimated costs resulting from incomplete adherence in the United States in 2001 were up to 300 billion dollars.¹² This figure highlights the importance of comprehensive knowledge about incomplete adherence to minimize the problems that are subsequently incurred by patients and health care systems worldwide.

Correspondence: Levente Kriston
 Department of Medical Psychology,
 University Medical Center Hamburg-
 Eppendorf Martinistraße 52
 20246, Hamburg, Germany
 Tel +49 40 7410 56849
 Fax +49 40 7410 54965
 Email l.kriston@uke.de

submit your manuscript | www.dovepress.com
 Dovepress
<http://dx.doi.org/10.2147/PPA.S38373>

Patient Preference and Adherence 2013:7 237-244

© 2013 Hüther et al, publisher and licensee Dove Medical Press Ltd. This is an Open Access article which permits unrestricted noncommercial use, provided the original work is properly cited.

237

Existing studies report various findings concerning the incomplete adherence of patients. Overall, incomplete adherence has been found to occur in 26% to 60% of all patients.^{13–15} For Germany, the results range between 35% and 50%,^{16,17} yet to this point, adherence has only been investigated in a few studies. In general, the findings vary depending on sample characteristics and applied measurement instruments.^{18,19} Concerning measurement, these different results may be due to the lack of a consistent method for accurate measurement of incomplete adherence.²⁰ Adherence measures include pill count, physical tests, medical records, self-report, collateral report, or electronic monitoring.¹² Concerning sample characteristics, the following health-related variables have been found to be statistically significantly associated with incomplete adherence: disease (particularly low subjective severity of the disease to be treated, or low “disease threat”;¹² also the occurrence of depression²¹); low objective severity of disease in patients with less serious conditions as well as a high objective severity of disease in patients with more serious conditions;²² low perceived need for medication;²³ lack of social support (including emotional support, family cohesiveness, and marital status);²⁴ poor communication between patient and physician;²⁵ high complexity of a patient’s medication regimen;^{16,18,26} and low extent of medication information (in this context, medication information means the level to which patients feel they have received enough information about prescribed medication).^{27,28} Unfortunately, these health-related factors are not always easy to assess in practice.

Demographic data are usually more feasible to acquire; however, according to a meta-analysis by DiMatteo,¹² demographic effects on adherence are small and moderated by sample, regimen, and measurement issues. The only sociodemographic factors found to be associated with incomplete adherence seem to be age below 50 and above 75 years,^{13,23} low income,^{12,23} and college education.²³ Yet, all these results were found to be affected by the population under study.¹² Most relevant studies have been conducted in the USA,²¹ whereas empirical evidence on incomplete medication adherence in Germany is still rare and should be amended by additional data.¹⁶ Accordingly, we aimed to assess the degree of incomplete medication adherence in a sample of chronically ill patients in German primary care. Furthermore, we examined if incomplete medication adherence is associated with certain health-related and/or sociodemographic characteristics. The following health-related variables were considered to potentially be associated with incomplete medication adherence: medication information,

medication complexity (consisting of dosage form, dosage frequency, and additional instructions), health-related quality of life (HRQoL, consisting of physical and mental health), and treated condition(s). Investigated sociodemographic data included age, sex, education, and employment status.

This analysis was conducted to clarify which of these potential associations should be taken into consideration in clinical care of chronically ill outpatients in German primary care. Once incomplete adherence is understood better, guidelines for screening procedures in clinical routine could be developed and ultimately lead to more favorable clinical outcomes and savings of health care costs.

Methods

Design

Data were collected in a prospective controlled trial that examined medication complexity, prescription behavior, and patient adherence at the interface between outpatient and inpatient medical care.²⁹ The data used for the presented analysis were obtained cross-sectionally at the time of admission during an inpatient stay for the treatment of at least one of the patient’s chronic cardiovascular and/or metabolic conditions. Patients were recruited between March 2010 and October 2011 from two internal medicine and two urology departments at the University Medical Center in Hamburg-Eppendorf, Germany. Patients were asked to assess their medication adherence in primary care treatment prior to admission retrospectively. Inclusion criteria for participation included age (>18 years), treatment for cardiovascular and/or metabolic diseases, sufficient knowledge of the German language, and the absence of cognitive impairment. To be included in the presented secondary analyses, a filled out German version of the Medication Adherence Report Scale (MARS-D) acquired from the patients was required to ensure that information on adherence was available.¹⁷

Measures

To measure incomplete medication adherence, the German version of the MARS-D was used.¹⁷ The MARS-D is considered to be an adequate tool for the detection of the frequency of patients’ incompletely adherent behavior. Internal consistency (Cronbach’s alpha: 0.60–0.69) and test–retest reliability (Pearson’s *r*: 0.61–0.63) of the MARS-D are satisfactory.¹⁷ The MARS-D consists of five items assessing the frequency of incompletely adherent behavior, each featuring a five-level Likert scale (from 1 = always to 5 = never) based on self-reports. Sum scores can vary between 5 and 25 points, with lower scores corresponding to incompletely

adherent behavior. There is no gold standard for dichotomizing the MARS-D and opinions differ concerning acceptable cut-off points,¹⁷ ranging from 20 to 25.^{31,32} In this study, patients were categorized as incompletely adherent if they scored less than 25 points on the MARS-D questionnaire. A high cut-off score (as used in this study) is recommended, as social desirability bias needs to be considered and any report of incomplete adherence should be taken into account.^{14,33} Additionally, we conducted sensitivity analyses with a cut-off of 23 (representing the lowest quartile of our distribution).

To measure medication information, the 17-item German version of the Satisfaction with Information about Medicines Scale (SIMS-D) was used.^{27,34} Total scores vary between 0 and 17 points, quantifying the patient's satisfaction with the information they received about their medication. The SIMS-D consists of two subscales: satisfaction with information received about medication usage (items 1–9) and satisfaction with information received about potential problems of the medication (items 10–17).

Medication complexity was obtained using the German version of the Medication Regimen Complexity Index (MRCI-D).^{7,35} The MRCI-D shows good interrater and test–retest reliability (intraclass correlation coefficients > 0.80) quantifying medication complexity concerning the total number of medications, the number of dosage units to be taken at a time, the dosage frequency, and specific directions concerning administration from clinical documentation.⁷

To measure HRQoL, the Short Form Health Survey Questionnaire-12, an indicator of physical and mental health, which was developed as a shorter version of the Short Form Health Survey Questionnaire-36, was utilized.^{36,37}

Patients also provided information on their sex, age, education, and employment status. Patients' diagnoses were acquired from their clinical records.

Statistical analyses

Descriptive analyses were used to describe the degree of incomplete adherence in this sample of chronically ill patients. Chi-squared/Fisher's exact tests were used to evaluate the relationship between categorical variables sex, education, employment status, diagnosis of hypertension/type two diabetes/hyperlipidemia/obesity/malignant tumor, and incomplete adherence, respectively. We used *t*-tests to assess differences between adherent and incompletely adherent patients regarding metric variables (age, medication information, medication complexity, HRQoL). To assess clinical significance, we report the respective effect sizes for all univariate results by calculating Cohen's *d* for all metric

and categorical variables. For the latter, we converted the respective odds ratios into Cohen's *d*.³⁸

To examine multivariate associations of sociodemographic and clinical information with incomplete adherence, we conducted a multiple logistic regression analysis. All variables were entered into the same model (entry method). In order to preclude collinearity problems, we entered total scores of instruments with highly correlating dimensions (SIMS-D, MRCI-D) instead of the subscales in the regression equation. The statistical analyses were performed with the software PASW/SPSS Statistics 18 (IBM Corp, Armonk, NY, USA).

Results

Descriptive analyses

A total of 190 patients met the inclusion criteria and took part in the study, providing an analysis sample of 142 male and 48 female participants. Of the 190 patients, 72 patients were recruited at urology departments. One hundred and fourteen patients were recruited at internal medicine departments. For four patients, this information was not available. Patients' age ranged from 23 to 92 years (mean = 62.9 years, standard deviation (SD) = 13.8). The mean MARS-D score was 23.5 (SD = 2.7). Forty-seven of the patients were employed, and 37 of them had an academic education (at least college). The patients' physical quality of life ranged from 15.8 to 61.5 (mean = 38.9, SD = 11.4). The patients' mental quality of life ranged from 18.2 to 64.2 (mean = 46.2, SD = 10.7). On average, patients reported being better informed about "action and usage of medication" (mean = 6.6, SD = 2.5) than about "potential problems of medication" (mean = 3.6, SD = 2.9). Overall, the patients scored an average 10.1 (SD = 4.8) points on the SIMS-D. Medication complexity was rather high with an average score of 15.2 (SD = 10.5) in the MRCI-D. Records of the patients' diseases were available for 186 of the initial 190 patients. The most common diseases were arterial hypertension (prevalence: 89.8%), type two diabetes (33.3%), malignant tumors (18.4%), obesity (13.4%), and hyperlipidemia (6.5%). In total, 62.1% (*n* = 118) of the patients were categorized as incompletely adherent.

Univariate analyses

Univariate analysis revealed no statistically significant differences at $P < 0.05$ between adherent and incompletely adherent patients regarding sex, age, employment status, academic education, medication information, medication complexity, physical and mental quality of life, hypertension, type two diabetes, obesity, or malignant tumors (Table 1).

Table 1 Associations between patient characteristics and medication adherence in univariate analysis

Variable	Complete adherence; N (%) 72 (37.9)	Incomplete adherence; N (%) 118 (62.1)	N (total) = 190	Test (df)	P	d
Sex; N (%)			190	$\chi^2 = 2.741$ (1)	0.098	0.307
Male	49 (25.8)	93 (48.9)				
Female	23 (12.1)	25 (13.2)				
Age; M (SD)	63.7 (14.8)	62.6 (13.2)	183	$t = -0.491$ (181)	0.624	0.08
Employment status; N (%)			189	$\chi^2 = 0.554$ (1)	0.457	0.144
Employed	16 (8.5)	31 (16.4)				
Unemployed	57 (30.2)	85 (44.9)				
Education; N (%)			183	$\chi^2 = 0.437$ (1)	0.508	0.140
Academic	13 (7.1)	24 (13.1)				
Not academic	60 (32.8)	86 (47)				
Patient information; M (SD)	11 (4.7)	9.6 (4.8)	178	$t = -1.922$ (176)	0.056	0.292
Action and usage	6.9 (2.4)	6.3 (2.5)	178	$t = -1.647$ (176)	0.101	
Potential problems	4 (2.9)	3.3 (2.9)	177	$t = -1.724$ (175)	0.086	
Medication complexity; M (SD)	16.5 (10.7)	14.4 (10.4)	175	$t = -1.301$ (173)	0.195	0.199
Physical QoL; M (SD)	39.6 (11.8)	38.6 (11.2)	174	$t = -0.541$ (172)	0.589	0.088
Mental QoL; M (SD)	46.2 (10.6)	46.1 (10.8)	174	$t = -0.056$ (172)	0.955	0.009
Hypertension; N (%)			186	$\chi^2 = 0.001$ (1)	0.981	0.007
Yes	62 (33.3)	105 (56.5)				
No	7 (3.8)	12 (6.4)				
DM type 2; N (%)			186	$\chi^2 = 0.104$ (1)	0.747	0.057
Yes	22 (11.8)	40 (21.5)				
No	47 (25.3)	77 (41.4)				
Hyperlipidemia; N (%)			186	FET	0.131	0.511
Yes	7 (3.8)	5 (2.7)				
No	62 (33.3)	112 (60.2)				
Obesity; N (%)			186	$\chi^2 = 2.123$ (1)	0.145	0.392
Yes	6 (3.2)	19 (10.2)				
No	63 (33.9)	98 (52.7)				
Malignant tumor; N (%)			190	$\chi^2 = 1.585$ (1)	0.208	0.282
Yes	10 (5.3)	25 (13.2)				
No	62 (32.6)	93 (48.9)				

Notes: Patient information is the patients' satisfaction with the information they received about their medication (SIMS-D score); medication complexity is the MRCI-D (Medication Regimen Complexity Index) Score.

Abbreviations: N, number; df, degrees of freedom; d, effect size; M, mean; SD, standard deviation; QoL, quality of life; FET, Fisher's exact test; DM Type 2, diabetes mellitus type two; SIMS-D, Satisfaction with Information about Medicines Scale; MRCI-D, Medication Regimen Complexity Index.

Conducting the identical univariate analysis using the alternative cut-off value (MARS-D score of 23) as part of the sensitivity analysis did not show any statistically significant results either. Most standardized effect sizes were negligible with a few, but still not statistically significant, moderate values.

Multivariate analyses

No strong multicollinearities between the variables were detected (Table 2); therefore, we conducted a multivariate analysis to determine if any of the variables were associated with incomplete medication adherence (Table 3). None of the variables were found to be statistically significantly associated with incomplete medication adherence. Conducting the identical multivariate analysis using the alternative cut-off

value (MARS-D score of 23) as part of the sensitivity analysis did not show any statistically significant results either. The total explained variance amounted to 11.8% (Nagelkerke's $R^2 = 0.118$).

Discussion

Due to the cross-sectional design of the study, the conducted analyses allowed us to investigate possible associations between incomplete medication adherence and the patients' sociodemographic and clinical data. However, cross-sectional studies are not appropriate for testing causal relationships between dependent and independent variables, and our findings should be interpreted accordingly. We found a rate of 62.1% of incomplete medication adherence in chronically ill patients. Neither univariate nor multivariate analysis showed

Table 2 Pairwise correlations between analyzed patient characteristics

	Sex	Age	Employment status	Education	SIMS-D score	MRCI-D score	Physical QoL	Mental QoL	Arterial hypertension	DM 2	Hyperlipidemia	Obesity	Malignant tumor
Sex	1												
Age	0.09	1											
Employment status	-0.12	-0.47*	1										
Education	-0.15*	0.00	0.12	1									
Patient information	-0.05	0.09	-0.06	0.01	1								
Medication complexity	0.19*	0.04	-0.14	-0.04	0.07	1							
Physical QoL	-0.08	-0.23*	0.24*	0.12	0.04	-0.29*	1						
Mental QoL	-0.12	0.02	-0.01	0.19*	0.17*	-0.26*	0.2*	1					
Arterial hypertension	0.08	0.07	-0.01	-0.01	-0.11	-0.04	0.06	0.01	1				
DM 2	-0.03	0.13	-0.03	-0.01	0.02	0.33*	-0.20*	0.04	0.17*	1			
Hyperlipidemia	0.17*	-0.01	0.01	0.04	0.02	0.07	-0.09	0.09	0.02	-0.04	1		
Obesity	-0.03	-0.10	0.04	-0.03	-0.13	-0.01	0.00	0.02	0.02	-0.15*	0.07	1	
Malignant tumor	-0.15*	0.15*	0.02	-0.05	-0.09	-0.23	0.07	-0.09	-0.09	-0.13*	0.07	0.07	1

Notes: *Level of significance < 0.05. Patient information represents the patients' satisfaction with the information they received about their medication (SIMS-D score). Medication complexity indicates the MRCI score. Abbreviations: R, Pearson product-moment correlation coefficient; SIMS-D, Satisfaction with Information about Medicines Scale; MRCI-D, Medication Regimen Complexity Index; QoL, quality of life; DM 2, diabetes mellitus type two.

Table 3 Associations of incomplete medication adherence in multivariate analysis

Variable	OR	95% CI	P
Female sex	1.38	0.55; 3.47	0.493
Lower age	0.99	0.96; 1.02	0.534
Unemployment	0.92	0.35; 2.40	0.859
Lower education	0.38	0.13; 1.11	0.076
Lower information (SIMS-D score)	0.94	0.87; 1.02	0.143
Lower medication complexity (MRCI-D score)	0.98	0.94; 1.02	0.211
Lower physical QoL	0.97	0.94; 1.01	0.108
Lower mental QoL	0.99	0.96; 1.03	0.756
Absence of arterial hypertension	0.56	0.17; 1.92	0.359
Absence of DM 2	0.78	0.31; 1.97	0.598
Hyperlipidemia	2.33	0.49; 11.11	0.288
Absence of obesity	0.82	0.28; 2.40	0.717
Absence of malignant tumor	0.96	0.35; 2.63	0.929

Note: Nagelkerke's $R^2 = 0.118$.

Abbreviations: OR, odds ratio; CI, confidence interval; SIMS-D, Satisfaction with Information about Medicines Scale; MRCI-D, Medication Regimen Complexity Index; QoL, quality of life; DM 2, diabetes mellitus type two.

evidence for significant associations between incomplete medication adherence and the patients' sociodemographic or clinical characteristics. Our findings conflict with preceding studies from other countries. Discrepancies between results of this study and other reported results regarding the associations of incomplete adherence could be due to several reasons, of which some include measurement of adherence, publication bias, low generalizability, and power.

First, a well-known problem when investigating incomplete medication adherence is measurement. A gold standard has not been developed, neither for the method of measurement, nor for the interpretation of results. Although direct measurements of medication ingestion, such as determining blood levels of pharmacological agents or medication event monitoring systems (a medication container with a special closure that records the time and date of each time the container is opened and closed), are more precise, questionnaires about the patient's adherence (self-reports) are usually used because they are cheaper, noninvasive, and easier to conduct. However, the adherence rates assessed through questionnaires depend on the patient's honesty and social desirability bias. While generally providing moderate-to-high concordance with objective measures, self-reports have been shown to provide higher adherence rates compared to nonself-reports (for example 13% higher than medication event monitoring systems, and 3% higher than pill counts).^{13,17,39,40} In this study the MARS-D (self-report) was used. While some researchers refer to the MARS as an appropriate measure to detect incomplete medication adherence,¹⁷ others do not recommend it, for it lacks sensitivity.³³ In order to increase sensitivity,

we decided to measure complete medication adherence (MARS-D score of 25) versus incomplete medication adherence (MARS-D score of less than 25). The measured incomplete adherence rate of 62.1% was rather high,¹² yet still in accordance with previous findings.^{15,32} However, sensitivity analyses with a broader MARS-D cut-off did not indicate any statistically significant associations between incomplete medication adherence and the patients' data. The findings suggest that the results are unlikely to be attributable to the strict cut-off that was chosen. However, additional investigation is needed to examine whether associations between adherence rates and patient characteristics depend on the type of measurement.

Secondly, a further explanation for contrary results could possibly be an underestimated publication bias in the field of adherence. Publication bias is a common problem in other fields of research.⁴¹ Since previous results on the associations of incomplete medication adherence have reported inconsistent findings with generally small effect sizes,²⁰ the extent of a possible publication bias should be explored through future meta-analyses. However, existing meta-analyses concluded that the possible risk of publication bias due to unpublished nonsignificant findings is rather low.^{21,24}

Thirdly, a limitation of our study is the convenient sample that was examined. However, participants were recruited consecutively without any selection, which makes the findings fairly generalizable to a population of patients admitted to inpatient treatment. This sample was heterogeneous, yet fairly representative of chronically ill patients treated with antihypertensive medication.²⁹ Differences regarding the patients' diseases in previous studies and this study could have influenced the results (and therefore limited this study), as (for example) depression and other diseases are shown to affect adherence.^{12,21} Nevertheless, multimorbidity is found in most samples of chronically ill patients and represents the same problem in research and practice. The majority (74.7%) of all participants were male; 38.7% of the patients were recruited from urology wards. However, since gender has not been found to influence adherence,¹² this should not reduce the generalizability of our findings. The relatively high average age of 62.9 years and the low percentage of patients being currently employed (24.9%) can be expected in a sample of chronically ill patients. Lastly, 79.8% of the participants were without an academic education. This finding is also expected in a sample of chronically ill patients because a low socioeconomic status is more likely to be accompanied by disease.^{42,43} The high clinical heterogeneity of the analyzed sample provides a meaningful, even if

negative, result on possible associations of adherence and medication. Based on the sufficient power of the analyses, it is probable that this negative finding contributes substantially to the existing knowledge by showing us that we are likely to know less than we sometimes suppose.

Finally, featuring a sample size of 190 patients, the power of this study allowed for the detection of any small-to medium-sized association (Cohen's *d* of 0.4) with a power above 80% (two-tailed at $P = 0.05$). Thus, it is to be expected that clinically relevant associations would have become visible. The total explained variance of 11.8% in the multivariate analysis indicates that even though combining information from all tested variables may explain a moderate amount of the observed variation in adherence behavior, the contribution of each single variable remains negligible.

In conclusion, we could not find sociodemographic or clinical variables that are associated with incomplete medication adherence. Variables associated with incomplete medication adherence found in previous research from other countries (and only one study regarding Germany) could not be confirmed. Further research is needed to find predictors of incomplete adherence in German settings. The focus should lie on primary care since inpatients exhibit almost complete medication adherence leading to more favorable clinical outcomes.¹³ Studies that explore medication adherence beyond individual patient characteristics are needed to fully understand adherence as an interactive construct shaped by the relationship between patients and their health care providers.⁴⁴

Disclosure

The authors report no conflicts of interest in this work.

References

1. Sicard P, Zeller M, Dentan G, et al; for RICO survey working group. Beneficial effects of statin therapy on survival in hypertensive patients with acute myocardial infarction: data from the RICO survey. *Am J Hypertens*. 2007;20(11):1133–1139.
2. Roik M, Starczewska MH, Huczek Z, Kochanowski J, Opolski G. Statin therapy and mortality among patients hospitalized with heart failure and preserved left ventricular function – a preliminary report. *Acta Cardiol*. 2008;63(6):683–692.
3. Halpern MT, Khan ZM, Schmier JK, et al. Recommendations for evaluating compliance and persistence with hypertension therapy using retrospective data. *Hypertension*. 2006;47(6):1039–1048.
4. DiMatteo MR, Giordani PJ, Lepper HS, Croghan TW. Patient adherence and medical treatment outcomes: a meta-analysis. *Med Care*. 2002;40(9):794–811.
5. Cramer JA, Benedict A, Muszbek N, Keskinaslan A, Khan ZM. The significance of compliance and persistence in the treatment of diabetes, hypertension and dyslipidaemia: a review. *Int J Clin Pract*. 2008;62(1):76–87.
6. Munger MA, Van Tassel BW, LaFleur J. Medication nonadherence: an unrecognized cardiovascular risk factor. *Med Gen Med*. 2007;9(3):58.

7. Stange D, Kriston L, Langebrake C, et al. Development and psychometric evaluation of the German version of the Medication Regimen Complexity Index (MRCI-D). *J Eval Clin Pract.* 2012;18(3):515–522.
8. McGinnis BD, Olson KL, Delate TM, Stolcpart RS. Statin adherence and mortality in patients enrolled in a secondary prevention program. *Am J Manag Care.* 2009;15(10):689–695.
9. Rasmussen JN, Chong A, Alter DA. Relationship between adherence to evidence-based pharmacotherapy and long-term mortality after acute myocardial infarction. *JAMA.* 2007;297(2):177–186.
10. Ho PM, Magid DJ, Masoudi FA, McClure DL, Rumsfeld JS. Adherence to cardioprotective medications and mortality among patients with diabetes and ischemic heart disease. *BMC Cardiovasc Disord.* 2006;6:48.
11. Sabate E. *Adherence to long-term therapies: evidence for action.* Geneva: World Health Organization; 2003.
12. DiMatteo MR. Variations in patients' adherence to medical recommendations: a quantitative review of 50 years of research. *Med Care.* 2004;42(3):200–209.
13. Dunbar-Jacob J, Mortimer-Stephens MK. Treatment adherence in chronic disease. *J Clin Epidemiol.* 2001;54 Suppl 1:S57–S60.
14. Haynes RB, McDonald HP, Garg AX. Helping patients follow prescribed treatment: clinical applications. *JAMA.* 2002;288(22):2880–2883.
15. Van Eijken M, Tsang S, Wensing M, de Smet PA, Grol RP. Interventions to improve medication compliance in older patients living in the community: a systematic review of the literature. *Drugs Aging.* 2003;20(3):229–240.
16. Wilke T, Müller S, Morisky DE. Toward identifying the causes and combinations of causes increasing the risks of nonadherence to medical regimens: combined results of two German self-report surveys. *Value Health.* 2011;14(8):1092–1100.
17. Mahler C, Hermann K, Horne R, et al. Assessing reported adherence to pharmacological treatment recommendations. Translation and evaluation of the Medication Adherence Report Scale (MARS) in Germany. *J Eval Clin Pract.* 2010;16(3):574–579.
18. Chapman RH, Benner JS, Petrilla AA, et al. Predictors of adherence with antihypertensive and lipid-lowering therapy. *Arch Intern Med.* 2005;165(10):1147–1152.
19. Dew MA, Dabbs AD, Myaskovsky L, et al. Meta-analysis of medical regimen adherence outcomes in pediatric solid organ transplantation. *Transplantation.* 2009;88(5):736–746.
20. Vermeire E, Hearnshaw H, Van Royen P, Denekens J. Patient adherence to treatment: three decades of research. A comprehensive review. *J Clin Pharm Ther.* 2001;26(5):331–342.
21. Grenard JL, Munjas BA, Adams JL, et al. Depression and medication adherence in the treatment of chronic diseases in the United States: a meta-analysis. *J Gen Intern Med.* 2011;26(10):1175–1182.
22. DiMatteo MR, Haskard KB, Williams SL. Health beliefs, disease severity, and patient adherence: a meta-analysis. *Med Care.* 2007;45(6):521–528.
23. Gadkari AS, McHorney CA. Unintentional non-adherence to chronic prescription medications: how unintentional is it really? *BMC Health Serv Res.* 2012;12(1):98.
24. DiMatteo MR. Social support and patient adherence to medical treatment: a meta-analysis. *Health Psychol.* 2004;23(2):207–218.
25. Zolnierak KB, DiMatteo MR. Physician communication and patient adherence to treatment: a meta-analysis. *Med Care.* 2009;47(8):826–834.
26. Schroeder K, Fahey T, Ebrahim S. How can we improve adherence to blood pressure-lowering medication in ambulatory care? Systematic review of randomized controlled trials. *Arch Intern Med.* 2004;164(7):722–732.
27. Mahler C, Jank S, Hermann K, et al. Psychometric properties of a German version of the "Satisfaction with Information about Medicines Scale" (SIMS-D). *Value Health.* 2009;12(8):1176–1179.
28. Gellaitry G, Cooper V, Davis C, Fisher M, Date HL, Horne R. Patients' perception of information about HAART: impact on treatment decisions. *AIDS Care.* 2005;17(3):367–376.
29. Stange D, Kriston L, von Wolff A, Baehr M, Dartsch DC. Medication complexity, prescription behaviour and patient adherence at the interface between ambulatory and stationary medical care. *Eur J Clin Pharmacol.* Epub July 25, 2012.
30. Ediger JP, Walker JR, Graff L, et al. Predictors of medication adherence in inflammatory bowel disease. *Am J Gastroenterol.* 2007;102(7):1417–1426.
31. George J, Kong DC, Thoman R, Stewart K. Factors associated with medication nonadherence in patients with COPD. *Chest.* 2005;128(5):3198–3204.
32. van de Steeg N, Sielk M, Pentzek M, Bakx C, Altiner A. Drug-adherence questionnaires not valid for patients taking blood-pressure-lowering drugs in a primary health care setting. *J Eval Clin Pract.* 2009;15(3):468–472.
33. Horne R, Hankins M, Jenkins R. The Satisfaction with Information about Medicines Scale (SIMS): a new measurement tool for audit and research. *Qual Health Care.* 2001;10(3):135–140.
34. George J, Phun YT, Bailey MJ, Kong DC, Stewart K. Development and validation of the medication regimen complexity index. *Ann Pharmacother.* 2004;38(9):1369–1376.
35. Brazier JE, Harper R, Jones NM, et al. Validating the SF-36 health survey questionnaire: new outcome measure for primary care. *BMJ.* 1992;305(6846):160–164.
36. Bullinger M. German translation and psychometric testing of the SF-36 Health Survey: preliminary results from the IQOLA Project. International Quality of Life Assessment. *Soc Sci Med.* 1995;41(10):1359–1366.
37. Cochrane-handbook.org [homepage on the Internet]. The Cochrane Collaboration, 2011. Cochrane Handbook for Systematic Reviews Interventions Version 5.1.0 [updated March 2011]. Available from: www.cochrane-handbook.org. Accessed February 9, 2013.
38. Horne R, Weinman J, Barber N, et al. Concordance, Adherence and Compliance in Medicine Taking. London: National Co-ordinating Centre for NHS Service Delivery and Organisation R and D; 2005. Available from: http://www.netscc.ac.uk/hsdr/files/project/SDO_FR_08-1412-076_V01.pdf. Accessed February 9, 2013.
39. Garber MC, Nau DP, Erickson SR, Aikens JE, Lawrence JB. The concordance of self-report with other measures of medication adherence: a summary of the literature. *Med Care.* 2004;42(7):649–652.
40. Robertson CT. The money blind: how to stop industry bias in biomedical science, without violating the First Amendment. *Am J Law Med.* 2011;37(23):358–387.
41. Kucharska-Newton AM, Harald K, Rosamond WD, Rose KM, Rea TD, Salomaa V. Socioeconomic indicators and the risk of acute coronary heart disease events: comparison of population-based data from the United States and Finland. *Ann Epidemiol.* 2011;21(8):572–579.
42. Secrest AM, Costacou T, Gutelius B, Miller RG, Songer TJ, Orchard TJ. Associations between socioeconomic status and major complications in type 1 diabetes: the Pittsburgh epidemiology of diabetes complication (EDC) Study. *Ann Epidemiol.* 2011;21(5):374–381.
43. Steiner JF. Rethinking adherence. *Ann Intern Med.* 2012;157(8):580–585.

SUMMARY

Introduction:

In all healthcare systems prescribing medication represents one of the cornerstones of medical treatment. The purpose of prescribing medications is to improve the patients' health, including the cure of disease.¹⁻⁵ The intake of an indicated medication can improve the patients' clinical outcome not only by treating the primary disease but also by reducing the probability of secondary diseases.⁶ Overall, patients who take their prescribed medication are less likely to be hospitalized or to die.⁷⁻¹⁰ In contrast, the complications resulting from patients not taking their medication accordingly are responsible for a big share of the costs in health care systems worldwide.¹¹

A patient's behavior of following a consensual treatment suggested by a health care provider can be referred to as *adherence*.¹² *Incomplete medication adherence* is therefore the occurrence of patients not completely following recommendations regarding their prescribed medication.

Current international literature suggests an average rate of incomplete adherence by between 26% and 60% of all patients.¹³⁻¹⁵ The average rate of adherence can vary quite heavily depending on sample and measurement,^{16,17} especially since there is a lack of a coherent method for the measurement of adherence.¹⁸ For Germany, the results range between 35% and 50%.^{19,20} However, the problem of incomplete adherence in Germany has only been examined by few studies.

Previous studies suggested several associations between incomplete adherence and patient characteristics, including socio-demographic data (low age,^{13,21} low income^{12,21} and college-education²¹) as well as clinical data

(disease,^{12,22} high complexity of a patient's medication regimen,^{16, 23} and low extent of medication information^{24,25}).

This dissertation explores the extent to which a sample of 190 German primary care patients suffering from chronic diseases adhere to their prescribed medication plan and if an association can be detected between the patients' medication adherence and their socio-demographic and clinical characteristics. Considering current literature, we rated the following health-related variables to be potentially associated with incomplete medication adherence: Sex, age, academic education, employment status, medication information (the level to which patients feel they have received enough information about their prescribed medication), medication complexity (consisting of dosage form, dosage frequency, additional instructions), health related quality of life (HRQoL - consisting of physical and mental health) and the prevalence of specific chronic diseases (hypertension, type-2 diabetes, hyperlipidemia, obesity, malignant tumor).

The analysis was conducted to clarify which of these potential associations should be taken into consideration in clinical care of chronically ill outpatients in German primary care. Once incomplete adherence is understood better, guidelines for screening procedures in clinical routine (e.g. via questionnaires) regarding the adherence of outpatients could be developed and ultimately lead to more favorable clinical outcomes and savings in health care costs.

Methods:

Data were collected in a prospective controlled trial that examined medication complexity, prescription behavior and patient adherence at the interface between inpatient and outpatient care.²⁶ The cross-sectional data used for the presented analysis were obtained at the time of admission as an inpatient

for the treatment of at least one of the patient's chronic cardiovascular and/or metabolic conditions. Patients were asked to assess their medication adherence in primary care treatment prior to admission retrospectively utilizing the German version of the Medication Adherence Report Scale (MARS-D).²⁰ Patients were categorized as incompletely adherent if they scored fewer than the maximum 25 points on the MARS-D questionnaire. A high cut-off score as used in this study is recommended, as social desirability bias is to be considered and any report of incomplete adherence should be taken into account.¹⁴ To ensure a high sensitivity, we additionally conducted all analyses mentioned below applying an alternative cut-off of 23 points (representing the lowest quartile of our distribution).

Descriptive analyses were used to describe the degree of incomplete adherence in this sample. Regarding the health-related variables (socio-demographic and clinical information that were rated to be potentially associated with incomplete medication adherence considering current literature) chi-squared/ Fisher's exact tests were used to evaluate the relationship between categorical variables (sex, education, employment status, diagnosis of hypertension/ type-2 diabetes/ hyperlipidemia/ obesity/ malignant tumor) and incomplete adherence. We used t-tests to assess differences between adherent and incompletely adherent patients regarding metric variables (age, medication information, medication complexity, HRQoL). To examine multivariate associations of socio-demographic and clinical information with incomplete adherence we conducted a multiple logistic regression analysis.




Figure 1. 1a) degree of adherence 1b) degree of medication information 1c) degree of medication complexity 2a) association of medication information and medication adherence 2b) association of medication complexity and medication adherence 2c) association of age, sex, education, employment status, quality of life, disease and medication adherence

Results:

A total of 190 patients met the inclusion criteria and took part in the study, providing an analysis sample of 142 male and 48 female participants. In total, 62.1% (n = 118) of the patients were categorized as incompletely adherent. None of the variables were found to be statistically significantly associated (at $p < 0.05$) with incomplete medication adherence in either univariate or multivariate analyses. This extends to the sensitivity analyses, using an alternative cut-off value of a MARS-D score of 23 points.

Discussion:

The results of this study indicate that the rates of incomplete medication adherence of German primary care patients with chronic disease are rather high (62.1%; measured via self-report). The cross-sectional design of this study only allowed testing for possible associations between the socio-demographic or clinical variables and incomplete medication adherence, a causal relationship was not to be detected. The finding that incomplete medication adherence is not associated with the patients' socio-demographic or clinical data conflicts with the results of preceding studies, the vast majority of them being from other countries. Possible reasons accounting for these conflicting results include the applied measurement of adherence, a possible publication bias in the field of medication adherence and a low generalizability of this sample.

To measure the patients' medication adherence, the MARS-D questionnaire was utilized. In the absence of a gold standard for the measurement of medication adherence, using different measurement tools can lead to variant results when testing for medication adherence.²⁷ Even though direct measurements of medication ingestion (determining blood levels of pharmacological agents) or medication event monitoring systems (MEMS = Medication container with a special closure that records the time and date of each time the container is opened and closed) are said to be more precise, questionnaires about the patient's adherence (self-reports) are usually used when investigating adherence because they are cheaper, non-invasive, and easier to conduct. However, the adherence rates assessed through questionnaires depend on the patient's honesty and social desirability bias. While generally providing moderate-to-high concordance with objective measures, self-reports have been shown to provide higher adherence rates compared to non-self-report (for example 13% higher than MEMS and 3%

higher than pill counts).^{13,20,27,28} Nevertheless, the MARS-D is considered an appropriate tool to measure medication adherence.²⁰ Although there is no determined cut-off value and therefore no guideline about how to interpret the results, by conducting sensitivity analyses the possibility that different cut-off values lead to different results was excluded. Additional investigations are needed to examine to which extent associations between adherence rates and patient characteristics depend on the type of measurement. The goal should be to find a gold-standard measurement tool that is easy to conduct in a primary care setting and detects incomplete medication adherence accurately.

A further explanation for the inconsistent results could be an underestimated publication bias in the field of adherence. Publication bias is a common problem in other fields of research.²⁹ A previous investigation on the associations of incomplete medication adherence by Vermeire et al.¹⁸ reported inconsistent findings with generally small effect sizes. The extent of a possible publication bias should be explored through future meta-analyses. However, meta-analyses that used fail-safe *n* have estimated the possible risk of unpublished non-significant results in the field of medication adherence to be rather low.^{22,30}

Another possible reason for the disparity of the results is a low generalizability. A limiting factor to the generalizability of this study's results is the convenience sample that was examined. Including 142 male and 48 female participants, the majority of this sample was male but since gender was not found to be associated with adherence,¹¹ the generalizability should not be reduced by the gender distribution. Differences regarding the patients' diseases in previous studies and this study could have influenced the results and therefore limited this study, as, for example, depression²² and other diseases¹² are shown to affect adherence. Nevertheless, multi-morbidity is found in most samples of chronically ill patients and

represents the same problem in research and practice. At least, the participants were recruited consecutively without selection, making up a heterogeneous and fairly representative sample of chronically ill patients treated with medication for chronic diseases like hypertension.²⁶ The high clinical heterogeneity of the analyzed sample provides a meaningful, albeit negative, result on the possible associations of adherence and medication. Based on the sufficient power of the analyses, this negative finding contributes substantially to existing knowledge by showing that we are likely to know less than we sometimes suppose.

The inconsistent results of previous studies investigating incomplete adherence and its associations suggest that incomplete adherence is not just a simple variable that is influenced by a few parameters. Adherence should maybe not be understood as a patient's characteristic, but rather as a complex and interactive construct that depends on several factors e.g. a patient's personality and the relationship between the patient and his/her health care provider.³¹ If the latter was to be confirmed in future studies, strategies to reduce incomplete medication adherence could include interventions focusing on the individual patient as well as interventions focusing on the patient-doctor relationship.

A very important precondition for better adherence is the patient's will to adhere to the prescribed medication. However, it has been shown that it is not sufficient simply to want something in order to achieve it.³² A person needs an efficient strategy to overcome the intention-behavior gap. A possible strategy to raise adherence rates is the administration of multi-focused interventions including cognitive, behavioral and affective components. Multi-focused interventions are more likely to raise rates of patients' medical adherence than single-focus interventions.³³ Strategies that have been shown to help people to attain their goals are, for instance, *Mental Contrasting* and *Implementation Intentions*.^{32,34,35} Future

studies applying such strategies are needed to confirm a positive effect on medication adherence.

Another crucial aspect that seems to influence a patient's adherence is the patient-doctor relationship¹² as poor communication between patients and their physicians can be associated with higher rates of incomplete adherence.³⁶ If the patient-doctor relationship is poor, the patient may not feel free to address his/her concerns regarding the prescribed medication. However, the patients' individual perceptions are crucial in their decision-making³⁷ and incomplete adherence, even though it may appear irrational to doctors and researchers, can represent a gain in quality of life for patients, for example by reducing side effects. Side effects are the highest concern patients have regarding their information about their medication.^{31,38} Therefore, specific training for doctors, teaching them how to address patients' risks and reasons not to adhere to their medication regimen properly, could improve adherence rates in primary care. Future research should examine the influence of such communication training for doctors on the medication adherence of their patients.

In conclusion, at this point we do not have sufficient knowledge about the reasons that account for incomplete medication adherence. Further investigation is needed to explore those reasons and how they can be addressed sufficiently in order to raise adherence rates and thus to make for more favorable clinical outcomes and savings in health care costs.

References:

1. Sicard P, Zeller M, Dentan G, et al. RICO survey working group: Beneficial effects of statin therapy on survival in hypertensive patients with acute myocardial infarction: data from the RICO survey. *Am J Hypertens* 2007, 20(11):1133-9.
2. Roik M, Starczewska MH, Huczek Z, et al. Statin therapy and mortality among patients hospitalized with heart failure and preserved left ventricular function - a preliminary report. *Acta Cardiol* 2008, 63(6):683-92.
3. Halpern MT, Khan ZM, Schmier JK, et al. Recommendations for evaluating compliance and persistence with hypertension therapy using retrospective data. *Hypertension* 2006, 47(6):1039-48.
4. DiMatteo MR, Giordani PJ, Lepper HS, et al. Patient adherence and medical treatment outcomes: a meta-analysis. *Med Care* 2002, 40(9):794-811.
5. Cramer JA, Benedict A, Muszbek N, et al. The significance of compliance and persistence in the treatment of diabetes, hypertension and dyslipidaemia: a review. *Int J Clin Pract* 2008, 62(1):76-87.
6. Munger MA, Van Tassell BW, LaFleur J: Medication nonadherence: an unrecognized cardiovascular risk factor. *MedGenMed* 2007, 9(3):58.
7. Stange D, Kriston L, Langebrake C, et al. Development and psychometric evaluation of the German version of the Medication Regimen Complexity Index (MRCI-D). *J Eval Clin Pract* 2012, 18(3):515-522.
8. McGinnis BD, Olson KL, Delate TM, et al. Statin adherence and mortality in patients enrolled in a secondary prevention program. *Am J Manag Care* 2009, 15(10):689-95.

9. Rasmussen JN, Chong A, Alter DA: Relationship between adherence to evidence-based pharmacotherapy and long-term mortality after acute myocardial infarction. *JAMA* 2007, 297(2):177-86.
10. Ho PM, Magid DJ, Masoudi FA, et al. Adherence to cardioprotective medications and mortality among patients with diabetes and ischemic heart disease. *BMC Cardiovasc Disord* 2006, 6:48.
11. DiMatteo MR: Variations in patients' adherence to medical recommendations: a quantitative review of 50 years of research. *Med Care* 2004, 42(3):200-9.
12. Sabate E: Adherence to long-term therapies: Evidence for action; 2003. Available from:
http://www.who.int/chp/knowledge/publications/adherence_report/en.
Accessed: February 9, 2013.
13. Dunbar-Jacob J, Mortimer-Stephens MK: Treatment adherence in chronic disease. *J Clin Epidemiol* 2001, 54(12):57-60
14. Haynes RB, McDonald HP, Garg AX: Helping patients follow prescribed treatment: clinical applications. *JAMA* 2002, 288(22):2880-3.
15. Van Eijken M, Tsang S, Wensing M, et al. Interventions to improve medication compliance in older patients living in the community: A systematic review of the literature. *Drugs Aging* 2003, 20(3):229-240.
16. Chapman RH, Benner JS, Petrilla AA, et al. Predictors of adherence with antihypertensive and lipid-lowering therapy. *Arch Intern Med* 2005, 165(10):1147-52.
17. Dew MA, Dabbs AD, Myaskovsky L, et al. Meta-analysis of medical regimen adherence outcomes in pediatric solid organ transplantation. *Transplantation* 2009, 88(5):736-46.

18. Vermeire E, Hearnshaw H, Van Royen P, et al. Patient adherence to treatment: three decades of research. A comprehensive review. *J Clin Pharm Ther* 2001, 26(5):331-342.
19. Wilke T, Müller S, Morisky DE: Toward identifying the causes and combinations of causes increasing the risks of nonadherence to medical regimens: combined results of two German self-report surveys. *Value Health* 2011, 14(8):1092-100.
20. Mahler C, Hermann K, Horne R, et al. Assessing reported adherence to pharmacological treatment recommendations. Translation and evaluation of the Medication Adherence Report Scale (MARS) in Germany. *J Eval Clin Pract* 2010, 16(3):574-579.
21. Gadkari AS, McHorney CA: Unintentional non-adherence to chronic prescription medications: How unintentional is it really? *BMC Health Serv Res* 2012, 12(1):98.
22. Grenard JL, Munjas BA, Adams JL, et al. Depression and medication adherence in the treatment of chronic diseases in the United States: a meta-analysis. *J Gen Intern Med* 2011, 26(10):1175-82.
23. Schroeder K, Fahey T, Ebrahim S: How Can We Improve Adherence to Blood Pressure-Lowering Medication in ambulatory Care? Systematic Review of Randomized Controlled Trials. *Arch Intern Med* 2004; 164(7):722-732.
24. Mahler C, Jank S, Hermann K, et al. Psychometric properties of a German version of the "satisfaction with information about medicines scale" (SIMS-D). *Value Health* 2009, 12(8):1176-1179.
25. Gellaitry G, Cooper V, Davis C, et al. Patients' perception of information about HAART: Impact on treatment decisions. *AIDS Care* 2005, 17(3):367-376.

26. Stange D, Kriston L, von Wolff A, et al. Medication complexity, prescription behaviour and patient adherence at the interface between ambulatory and stationary medical care. *Eur J Clin Pharmacol* 2013, 69(3):573-80.
27. Horne R, Weinman J, Barber N, et al. Concordance, Adherence and Compliance in Medicine Taking. London: National Co-ordinating Centre for NHS Service Delivery and Organisation R and D; 2005. Available from:
http://www.netscc.ac.uk/hsdr/files/project/SDO_FR_08-1412-076_V01.pdf.
Accessed: February 9, 2013.
28. Garber MC, Nau DP, Erickson SR, et al. The concordance of self-report with other measures of medication adherence: a summary of the literature. *Med Care* 2004, 42(7):649-52.
29. Robertson CT: The money blind: how to stop industry bias in biomedical science, without violating the First Amendment. *Am J Law Med* 2011, 37(23):358-87.
30. DiMatteo MR: Social support and patient adherence to medical treatment: a meta-analysis. *Health Psychol* 2004, 23(2):207-18.
31. Steiner JF: Rethinking Adherence. *Ann Intern Med* 2012, 157:580-585.
32. Gollwitzer PM: Goal achievement: The role of intentions. *Eur Rev Sol Psychol* 1993, 4(1):141-185.
33. Roter DL, Hall JA, Merisca R: Effectiveness of Interventions to Improve Patient Compliance: A Meta-Analysis. *Med Care* 1998; 36(8):1138-61.
34. Gollwitzer PM: Implementation intentions: Strong effects of simple plans. *Am Psychol* 1999, 54(7):493-503.
35. Oettingen G: Future thought and behavior change. *Eur Rev Soc Psychol* 2012, 23(1):1-63.

36. Zolnieriek KB, Dimatteo MR: Physician communication and patient adherence to treatment: a meta-analysis. *Med Care* 2009, 47(8):826-34.
37. Donovan JL, Blake DR: Patient non-compliance: deviance or reasoned decision-making? *Soc Sci Med* 1992; 34(5):507-13.

AUTHOR CONTRIBUTION

Jakob Hüther conceptualized the aim and the design of the study, managed the data, performed the statistical analyses, participated in the primary interpretation of findings and drafted the manuscript.

CO-AUTHOR CONTRIBUTION

Alessa von Wolff participated in the conception and design of the study, performed the statistical analyses, participated in the primary interpretation of findings and revised the manuscript critically for important intellectual content.

Dorit Stange participated in the conception and design of the study, collected the data, participated in the primary interpretation of findings and revised the manuscript critically for important intellectual content.

Levente Kriston participated in the conception and design of the study, participated in the primary interpretation of findings and revised the manuscript critically for important intellectual content.

Michael Baehr participated in the conception and design of the study and revised the manuscript critically for important intellectual content.

Dorothee C Dartsch participated in the conception and design of the study and revised the manuscript critically for important intellectual content.

Martin Härter participated in the conception and design of the study and revised the manuscript critically for important intellectual content.

All authors read and approved the final manuscript.

ACKNOWLEDGEMENTS

This research project, and therefore this dissertation, would not have been possible without the support of the author's doctorate supervisor Professor Dr. Dr. Martin Härter who lead and oversaw the process of this dissertation with extraordinary professional knowledge, experience and guidance.

My deepest gratitude is also due to the supervisory committee, Dr. phil. Levente Kriston and Dipl. Psych. Alessa von Wolff without whose contagious excitement, profound knowledge and invaluable advice this dissertation would have never reached completion.

Further, I deeply thank Dorit Stange, Dr. Michael Baehr and Prof. Dr. Dorothee C. Dartsch for supporting me with great care, always providing important intellectual content.

Special thanks are due to Bettina Schwörer for incredible patience and priceless assistance.

The author would also like to convey great appreciation to the Department of Medical Psychology at the University Medical Center Hamburg-Eppendorf in Hamburg, Germany for providing the financial means.

Finally, the author wishes to express his love and gratitude to his beloved family for their understanding, support and endless love.

- Seite entfällt aus datenschutzrechtlichen Gründen -

EIDESSTATTLICHE VERSICHERUNG

Ich versichere ausdrücklich, dass ich die Arbeit selbständig und ohne fremde Hilfe verfasst, andere als die von mir angegebenen Quellen und Hilfsmittel nicht benutzt und die aus den benutzten Werken wörtlich oder inhaltlich entnommenen Stellen einzeln nach Ausgabe (Auflage und Jahr des Erscheinens), Band und Seite des benutzten Werkes kenntlich gemacht habe.

Ferner versichere ich, dass ich die Dissertation bisher nicht einem Fachvertreter an einer anderen Hochschule zur Überprüfung vorgelegt oder mich anderweitig um Zulassung zur Promotion beworben habe.

Ich erkläre mich einverstanden, dass meine Dissertation vom Dekanat der Medizinischen Fakultät mit einer gängigen Software zur Erkennung von Plagiaten überprüft werden kann.

Unterschrift: J. Höcher